Allergy Action Plan

THIS CHILD HAS THE FOLLOWING ALLERGIES:

Name:

DOB:

Photo

Emergency contact details:

1) ☎

2) ☎

Child's Weight: Kg


How to give EpiPen®




Form fist around EpiPen® and PULL OFF BLUE SAFETY CAP



SWING AND PUSH ORANGE TIP against outer thigh (with or without clothing) until a click is heard



HOLD FIRMLY in place for 10 seconds



REMOVE EpiPen®. Massage injection site for 10 seconds

Keep your EpiPen device(s) at room temperature, do not refrigerate.
For more information and to register for a free reminder alert service, go to www.epipen.co.uk

Produced in conjunction with:

Allergy UK – leading the fight against allergy

Anaphylaxis Campaign – Supporting people with severe allergies

©The British Society for Allergy & Clinical Immunology
www.bsaci.org Approved Oct 2013

Mild-moderate allergic reaction:

- Swollen lips, face or eyes
- Itchy / tingling mouth
- Hives or itchy skin rash
- Abdominal pain or vomiting
- Sudden change in behaviour

ACTION:

- Stay with the child, call for help if necessary
- Give antihistamine: ▼
- Contact parent/carer

(if vomited, can repeat dose)

Watch for signs of ANAPHYLAXIS
(life-threatening allergic reaction):

AIRWAY:	Persistent cough, hoarse voice, difficulty swallowing, swollen tongue
BREATHING:	Difficult or noisy breathing, wheeze or persistent cough
CONSCIOUSNESS:	Persistent dizziness / pale or floppy suddenly sleepy, collapse, unconscious

If ANY ONE of these signs are present:

1. **Lie child flat.** If breathing is difficult, allow to sit
2. **Give EpiPen® or EpiPen® Junior**
3. **Dial 999 for an ambulance*** and say ANAPHYLAXIS ("ANA-FIL-AX-IS")

If in doubt, give EpiPen®

After giving Epipen:

1. Stay with child, contact parent/carer
2. Commence CPR if there are no signs of life
3. If no improvement **after 5 minutes, give a further EpiPen®** or alternative adrenaline autoinjector device, if available

*You can dial 999 from any phone, even if there is no credit left on a mobile.
Medical observation in hospital is recommended after anaphylaxis.

Additional instructions:

If wheezy, give 10 puffs salbutamol (blue inhaler) via spacer an ▼

This is a medical document that can only be completed by the patient's treating health professional and cannot be altered without their permission.

This plan has been prepared by: _____

Hospital/Clinic:

☎ Date: