


My Asthma Plan



Your asthma plan tells you when to take your asthma medicines.

And what to do when your asthma gets worse.



Name: _____

1 My daily asthma medicines

- My preventer inhaler is called _____ and its colour is _____
- I take _____ puff/s of my preventer inhaler in the morning and _____ puff/s at night. I do this every day even if I feel well.
- Other asthma medicines I take every day:

- My reliever inhaler is called _____ and its colour is _____.
I take ____ puff/s of my reliever inhaler (usually blue) when I wheeze or cough, my chest hurts or it's hard to breathe.
- My best peak flow is _____

2 When my asthma gets worse

I'll know my asthma is getting worse if:

- I wheeze or cough, my chest hurts or it's hard to breathe, or
- I'm waking up at night because of my asthma, or
- I'm taking my reliever inhaler (usually blue) more than three times a week, or
- My peak flow is less than _____

If my asthma gets worse, I should:


Keep taking my preventer medicines as normal.

And also take _____ puff/s of my blue reliever inhaler every four hours.



If I'm not getting any better doing this I should see my doctor or asthma nurse today.


Does doing sport make it hard to breathe?



If YES

I take:

_____ puff/s of my reliever inhaler (usually blue) beforehand.



Remember to use my inhaler with a spacer (if I have one)

My Asthma Plan

3 When I have an asthma attack

I'm having an asthma attack if:

- My blue reliever inhaler isn't helping, or
- I can't talk or walk easily, or
- I'm breathing hard and fast, or
- I'm coughing or wheezing a lot, or
- My peak flow is less than _____

When I have an asthma attack, I should:

Sit up — don't lie down. Try to be calm.

Take one puff of my reliever inhaler **every 30 to 60 seconds** up to a total of 10 puffs.

Even if I start to feel better, I don't want this to happen again, so I need to see my doctor or asthma nurse today.



If I still don't feel better and I've taken ten puffs, I need to call **999** straight away. If I am waiting longer than 15 minutes for an ambulance I should take another _____ puff/s of my blue reliever inhaler every 30 to 60 seconds (up to 10 puffs).



My asthma triggers:


Write down things that make your asthma worse

I need to see my asthma nurse every six months

Date I got my asthma plan:

Date of my next asthma review:

Doctor/asthma nurse contact details:



Make sure you have your reliever inhaler (usually blue) with you. You might need it if you come into contact with things that make your asthma worse.

Parents – get the most from your child's action plan

Make it easy for you and your family to find it when you need it

- Take a **photo** and keep it on your mobile (and your child's mobile if they have one)
- Stick a **copy** on your fridge door
- Share your child's action plan with school, grandparents and babysitter (a printout or a photo).

You and your parents can get your questions answered:

Call our friendly expert nurses

0300 222 5800

(9am – 5pm; Mon – Fri)

Get information, tips and ideas

www.asthma.org.uk